

Generalizing a Closed-Form Correlation Model of Oriented Bandpass Natural Images

ZEINA SINNO – ALAN BOVIK

THE IEEE GLOBAL CONFERENCE ON SIGNAL AND INFORMATION PROCESSING
(GLOBALSIP)


14/12/2015

Outline

- Background
- The Basic Image Model
- The Generalized Model
- Validation of the Model
- Conclusion and Future Work

Photos to the Sun and Back


- **One trillion pictures** were captured in 2015.
- If the stored photos in a year are printed as 4x6 inches they would stretch 1.1 roundtrips from the Earth to the Sun.
- In 2017, the trail will stretch to make 2.5 roundtrips. ^[1]



[1]: Infotrends 2014 Worldwide Image Capture Forecast.

Photos to the Sun and Back

- **One trillion pictures** were captured in 2015.
- If the stored photos in a year are printed as 4x6 inches they would stretch 1.1 roundtrips from the Earth to the Sun.
- In 2017, the trail will stretch to make 2.5 roundtrips. ^[1]




➔ Understanding the **Natural Scene Statistics** is of a great importance.

[1]: Infotrends 2014 Worldwide Image Capture Forecast.

Natural Scene Statistics (NSS)

- Ruderman applied a simple form of “predictive coding” filter to a large numbers of images and observed that images follow an empirical probability distribution function.
- NSS follow **well behaved** and **universal models**.
- Combining NSS with Human Visual System provides **powerful tools**:
 - Image Denoising [2]
 - Image and Video Quality Assessment [3]
 - Image Defocus [4]
 - ...



[1]: D.L., “The Statistics of Natural Images, *Nature*, 1994.

[2]: J. Perry and W. S. Geisler, “Natural scene statistics for image denoising,” Center for Perceptual Systems, The University of Texas at Austin, Tech. Rep., 2013

[3]: Z. Wang, A. C. Bovik, H. Sheikh, and E. Simoncelli, “Image quality assessment: From error visibility to structural similarity,” *IEEE Trans. on Imag. Proc.*, vol. 13, no. 4, pp. 600–612, 2004.

[4]: J. Burge and W. S. Geisler, “Optimal defocus estimation in individual natural images,” *Proc. of the Nat. Acad. of Sci.*, vol. 108, no. 40, pp. 16 849–16 854, 2011

Natural Scene Statistics (NSS)

Extending NSS to understand the relationship between two pixels, referred as Bivariate NSS and more pixels is **useful** for:

- Understanding correlated noise → better denoising algorithms
- Estimating the depth from single images
- Modeling textures
- Estimating the quality of images more closely to human judgement
- ...

Background

- Previous work on NSS focused mainly on Univariate NSS.
- Random processes were used to model Bivariate NSS.

 **No closed-form models!**

- Su *et. al* ^[1] modeled spatially adjacent bandpass image responses.
- Model proved to be useful for:
 - Depth Estimation from Single Images ^[2]
 - Stereoscopic Image quality Prediction ^[3]

Goal: To extend this model over more pixels and more scales.

[1]: C. C. Su, L. Cormack, and A. C. Bovik, "Closed-form correlation model of oriented bandpass natural images," *Signal Processing Lett. IEEE*, vol. 22, no. 1, pp. 21–25, Jan 2015.


[2]: C. C. Su, L. K. Cormack, and A. C. Bovik, "Bivariate statistical modeling of color and range in natural scenes," in Proc. SPIE, *Human Vis. Electron. Imag. XIX*, vol. 9014, Feb. 2014.

[3]: C. Su, L. Cormack, and A. Bovik, "Oriented correlation models of distorted natural images with application to natural stereopair quality evaluation," *IEEE Trans. Image Process.*, vol. 24, no. 5, pp. 1685–1699, May 2015.

The Basic Image Model

Model constructed using 100 pristine images:

- 29 images from the LIVE Database ^[1]
- 71 images from the Berkeley Segmentation Database ^[2]



[1]: H. Sheikh, M. Sabir, and A. Bovik, "A statistical evaluation of recent full reference image quality assessment algorithms," *IEEE Trans. Image Processing*, vol. 15, no. 11, pp. 3440–3451, Nov 2006

[2]: D. Martin, C. Fowlkes, D. Tal, and J. Malik, "A database of human segmented natural images and its application to evaluating segmentation algorithms and measuring ecological statistics," in *Int. Conf. Comput. Vision*, vol. 2. IEEE, 2001, pp. 416–423.


The Basic Image Model

- Luminance images considered as they relate better to the human perception of colors.
- Steerable filters were applied as a simple model of simple cells in the primary visual cortex (area V1).
- Steerable filters at an orientation θ_1 :

$$F(\theta_1) = \cos(\theta_1)F_x + \sin(\theta_1)F_y$$

F_x and F_y : Gradient of a 2D Gaussian of variance σ w.r.t to the x and y axis.

- Considered 15 different θ_1 spanning $[0, \pi)$ with steps of $\frac{\pi}{15}$.



Steerable Filtering on Luminance Images



Divisive Normalization



Modeling the Bivariate Joint Distribution



Modeling the Correlation between Pixels

The Basic Image Model

- Divisive Normalization was applied to model the non-linear adaptive gain control of neuronal responses in the area V1.

- It is represented as:

$$u(x_i, y_i) = \frac{w(x_i, y_i)}{\sqrt{s + \sum_j g(x_j, y_j)w(x_j, y_j)^2}}$$


(x_i, y_i) : spatial coordinates

$u(x_i, y_i)$: coefficient after divisive normalization

$g(x_j, y_j)$: Gaussian weighting function

$w(x_i, y_i)$: steerable filters coefficients

s : stabilization constant



Steerable Filtering on Luminance Images


Divisive Normalization

Modeling the Bivariate Joint Distribution

Modeling the Correlation between Pixels

The Basic Image Model

- The bivariate joint distribution characterizes pixels between distance 1 to 10, at a spatial orientation θ_2 .
- Define the relative orientation as: $\theta_2 - \theta_1$



Steerable Filtering on Luminance Images

Divisive Normalization

Modeling the Bivariate Joint Distribution

Modeling the Correlation between Pixels

The Basic Image Model

- This is modeled using a Bivariate Generalized Gaussian Distribution (BGGD):


$$p(\mathbf{x}; \mathbf{M}, \alpha, \beta) = \frac{1}{|\mathbf{M}|^{\frac{1}{2}}} g_{\alpha, \beta}(\mathbf{x}^T \mathbf{M}^{-1} \mathbf{x})$$

\mathbf{x} : Pixels to be correlated

\mathbf{M} : Scatter Matrix


α, β : Shape Parameters

$g_{\alpha, \beta}$: Density generator function




The Generalized Image Model

- Take the correlation between the pixels after the DN for:
 - a spatial separation in $[0,10]$
 - at all possible digital angles
 - for scales σ in $[1,4]$ with steps of 0.5 (7 scales in total)
- Su *et. al* ^[1] considered horizontally adjacent pixels only and 5 scales.



The Generalized Image Model


- Su *et. al* 's Model [1]



[1]: C. C. Su, L. Cormack, and A. C. Bovik, "Closed-form correlation model of oriented bandpass natural images," *Signal Processing Lett. IEEE*, vol. 22, no. 1, pp. 21–25, Jan 2015.

The Generalized Image Model

- Su *et. al* 's Model [1] 



A
More
General
Model

[1]: C. C. Su, L. Cormack, and A. C. Bovik, "Closed-form correlation model of oriented bandpass natural images," *Signal Processing Lett. IEEE*, vol. 22, no. 1, pp. 21–25, Jan 2015.

The Generalized Model

- Model the correlation as a function of $\theta_2 - \theta_1$:

$$\rho = A \cos(\theta_2 - \theta_1) + c$$

versus Su *et. al*'s model^[1] :

$$\rho = A \cos(\theta_2 - \theta_1)^{2\gamma} + c$$


The Generalized Model

- Model the correlation as a function of $\theta_2 - \theta_1$:

$$\rho = A \cos(\theta_2 - \theta_1) + c$$

versus Su *et. al*'s model^[1] :


$$\rho = A \cos(\theta_2 - \theta_1)^{2\gamma} + c$$



The Generalized Model



- Consistent model over all the scales and distances:

For $\sigma = 2.5$ and $\theta_2 = \frac{\pi}{2}$:



The Generalized Model

- Amplitude (A) and Offset (c) over multiple scales:



Validation of the Model

Model validated using Mean Squared Error (MSE) and Pearson's χ^2 test:

	MSE training	χ^2 training	MSE testing	χ^2 testing
Distance 1	0.004178	0.459442	0.001053	0.11582
Distance 3	0.166615	48.77902	0.038803	11.43254
Distance 5	0.364689	-27.6250	0.070773	-0.46758
Distance 7	0.323636	290.3075	0.057003	47.40307

Training Data: the 100 images used to build the model

Testing Data: the 23 pristine images of the VCL@FER database^[1]

[1]: A. Zaric, N. Tatalovic, N. Brajkovic, H. Hlevnjak, M. Loncaric, E. Dumic, and S. Grgic, "Vcl@ fer image quality assessment database," *AUTOMATIKA*, vol. 53, no. 4, pp. 344–354, 2012.

Future Work

- Find a closed form for the amplitude and offset.
- Extend the model over more scales.
- Apply the model to problems including:
 - texture modeling
 - image quality prediction
 - image interpolation
 - ...

Thank you



Backup Slide

Density Generator Function:

$$g_{\alpha,\beta}(y) = \frac{\beta \Gamma(\frac{N}{2})}{(2^{\frac{1}{\beta}} \pi \alpha)^{\frac{N}{2}} \Gamma(\frac{N}{2\beta})} e^{-\frac{1}{2}(\frac{y}{\alpha})^{\beta}}$$

Γ : the Digamma function

$y \in \mathbb{R}^+$

α, β : Shape Parameters